

ON THE MAXIMAL DISTANCE BETWEEN TWO RENEWAL EPOCHS

P. RÉVÉSZ

*Institute for Statistics and Probability, Technical University, Vienna, Austria and Mathematical Institute, Budapest, Hungary**

E. WILLEKENS

Department of Mathematics, Catholic University of Leuven, Belgium

Received 13 March 1987

Revised 9 July 1987

Let X_1, X_2, \dots be a sequence of positive, independent, identically distributed (i.i.d.) random variables with $S_0 = 0$, $S_n = X_1 + \dots + X_n$, $n \geq 1$. Denote by $\tau_t = \sup\{n \mid S_n \leq t\}$. In this paper we establish almost sure lower and upper bounds for $M_t = \max\{X_1, X_2, \dots, X_{\tau_t}, t - S_{\tau_t}\}$ if the underlying distribution function has a regularly varying tail.

AMS (1980) Subject Classifications: 60F15, 60K05.

renewal processes * almost sure limit laws

1. Introduction

Let X_1, X_2, \dots be a sequence of positive i.i.d. random variables with $P(X_1 \leq x) = F(x)$. Further let $S_0 = 0$, $S_n = X_1 + \dots + X_n$, $n \geq 1$ and let τ_t be the largest integer for which $S_{\tau_t} \leq t$. The main goal of this paper is to study the properties of the process

$$M_t = M(t) = \max\{X_1, X_2, \dots, X_{\tau_t}, t - S_{\tau_t}\}.$$

M_t can be interpreted in its proper context, for example in a queueing process, it might represent the longest interval up to t without arrival of a customer.

The present problem is originated by a problem in random walk theory: in fact, let R_1, R_2, \dots be a sequence of i.i.d. random variables with $P(R_1 = 1) = \frac{1}{2} = P(R_1 = -1)$ and put

$$Y_0 = 0, \quad Y_n = R_1 + R_2 + \dots + R_n \quad (n \geq 1),$$

$$\rho_0 = 0, \quad \rho_1 = \min\{i > 0; Y_i = 0\},$$

$$\rho_{j+1} = \min\{i > \rho_j; Y_i = 0\} \quad (j \geq 1).$$

* Research done while visiting the Catholic University of Leuven, Belgium.

Let μ_t be the largest integer for which $\rho_{\mu_t} \leq t$. Then the length of the largest (not surely completed) excursion of the random walk $\{Y_n\}$ is given by

$$\varepsilon_t = \max\{\rho_1, \rho_2 - \rho_1, \dots, \rho_{\mu_t} - \rho_{\mu_t-1}, t - \rho_{\mu_t}\}.$$

It is clear that the problem of studying the properties of ε_t is a special case of the problem mentioned above. In fact, if $X_i = \rho_i - \rho_{i-1}$, then $M_t = \varepsilon_t$. We now summarize some of the results describing the behaviour of ε_t as $t \rightarrow \infty$. We start with a result of Chung and Erdős [4], giving the best possible upper bound for ε_t .

Theorem A [4; 5, p. 366]. *Let f be a non-decreasing positive function for which $\lim_{x \rightarrow \infty} f(x) = \infty$ and put*

$$I(f) = \int_1^\infty \frac{dx}{xf^{1/2}(x)}.$$

Then

$$P\left\{\varepsilon_n \geq n\left(1 - \frac{1}{f(n)}\right) \text{ i.o.}\right\} = \begin{cases} 1 & \text{if } I(f) = \infty, \\ 0 & \text{if } I(f) < \infty. \end{cases} \quad (1.1)$$

The best possible lower bound for ε_n was established by Csáki et al. [5]. They showed that

$$\liminf_{n \rightarrow \infty} \varepsilon_n \frac{\log \log n}{n} = \beta \quad \text{a.s.} \quad (1.2)$$

where β is the root of the equation

$$\sum_{k=1}^{\infty} \frac{\beta^k}{k!(2k-1)} = 1.$$

(1.1) implies that

$$\varepsilon_n \geq n - \frac{n}{(\log n)^2}$$

infinitely often with probability one. This shows that for some n , the random walk $\{Y_n\}$ is essentially one excursion. On the other hand, (1.2) shows that for some n , the length of the longest excursion is only $\beta n (\log \log n)^{-1}$, which implies that in these cases the random walk consists of at least $\beta^{-1} \log \log n$ excursions. The main theorems of the present paper extend the results in (1.1) and (1.2) to M_t .

Throughout the paper we assume that the underlying distribution function (d.f.) F has a regularly varying tail, i.e.

$$1 - F(x) = \frac{x^{-\alpha}}{L(x)}, \quad x > 0, \quad \alpha \geq 0, \quad (1.3)$$

with $L: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ slowly varying ($L \in SV$), which means that L is measurable, positive and

$$\lim_{x \rightarrow \infty} \frac{L(xt)}{L(x)} = 1 \quad \text{for every } t > 0.$$

The number $-\alpha$ in (1.3) is called the index of regular variation of $1 - F$.

Remark. In case of the random walk $\{Y_n\}$, we have that

$$P(\rho_1 = 2k) = \binom{2k-2}{k-1} \frac{1}{2^{2k-1} \cdot k} \approx \frac{k^{-3/2}}{2\sqrt{\pi}} \quad (k \rightarrow \infty)$$

which means that ε_n is just a special version of M_t when the underlying d.f. F has a regularly varying tail of index $-\frac{1}{2}$.

In the sequel we write

$$L_\alpha(x) := (L^{1/\alpha})^*(x^{1/\alpha}), \quad x > 0, \alpha > 0, \quad (1.4)$$

where L^* denotes the so called conjugate slowly varying function to L , see [3, 8]. With the definition in (1.4), it follows from de Bruyn [8] that

$$x^\alpha L(x) \sim y(x \rightarrow \infty)$$

if and only if

$$x \sim y^{1/\alpha} L_\alpha(y) \quad (y \rightarrow \infty),$$

To formulate our results, we also need a refined version of slow variation, provided by the so called π -variation (πV): $L: \mathbb{R}^+ \rightarrow \mathbb{R}$ is π -varying ($L \in \pi V$) if and only if there exists a function $a \in SV$ such that

$$\lim_{x \rightarrow \infty} \frac{L(xt) - L(x)}{a(x)} = \log t, \quad t > 0. \quad (1.5)$$

If (1.5) holds, de Haan [9] showed that it is always possible to take

$$a(x) = C(x) := L(x) - \frac{1}{x} \int_0^x L(y) dy. \quad (1.6)$$

The concept of π -variation was in its turn refined by Omey and Willekens [15]. They studied functions L satisfying

$$\lim_{x \rightarrow \infty} \frac{L(xt) - L(x) - a(x) \log t}{b(x)} = \Lambda(t) \quad (1.7)$$

for some functions a , b and Λ .

If L satisfies (1.7), we call it π -varying with remainder ($L \in \pi R$). It was shown in [15] that (1.7) holds with $b \in SV$ if and only if

$$\lim_{x \rightarrow \infty} \frac{C(xt) - C(x)}{b(x)} = \log t, \quad t > 0, \quad (1.8)$$

where C is the function defined in (1.6).

Furthermore if (1.7) holds, we can always take $a(x) = C(x)$ and $\Lambda(t) = \log t + \frac{1}{2}(\log t)^2$, see [15].

We are now ready to formulate our main results, presented in the next section. The proofs are collected in Section 3.

2. Main results

We distinguish between the two cases $\mu := \int_0^\infty x \, dF(x) < \infty$ and $\mu = \infty$. First consider the case $\mu < \infty$. It then follows from the strong law of large numbers that

$$\frac{\tau_t}{t} \rightarrow \frac{1}{\mu} \quad \text{a.s.}$$

Hence M_t can be approximated by

$$U_t := \max\{X_1, X_2, \dots, X_{[t\mu^{-1}]}\}$$

and both M_t and U_t have the same a.s. asymptotic behaviour. Best possible results on the a.s. limit behaviour of U_t were obtained by Galambos [11, Theorem 4.3.1, Corollary 4.3.1]. For the sake of completeness we translate his results to M_t :

Theorem 1. *Let $1 - F(x) = x^{-\alpha} / L(x)$ and suppose that $\mu < \infty$.*

Let g be a positive increasing function with $g(t) \uparrow \infty$ as $t \rightarrow \infty$, and define

$$I(g) = \int_1^\infty \frac{dt}{g(t)} \quad J(g) = \int_1^\infty \exp\left(\frac{x}{g(x)}\right) \frac{dx}{g(x)}.$$

Then

$$P\{M_t \geq (g(t/\mu))^{1/\alpha} L_\alpha(g(t/\mu)) \text{ i.o.}\} = \begin{cases} 1 & \text{if } I(g) = \infty, \\ 0 & \text{if } I(g) < \infty. \end{cases}$$

If furthermore $t/g(t)$ is nondecreasing as $t \rightarrow \infty$, then

$$P\{M_t \leq (g(t/\mu))^{1/\alpha} L_\alpha(g(t/\mu)) \text{ i.o.}\} = \begin{cases} 1 & \text{if } J(g) = \infty, \\ 0 & \text{if } J(g) < \infty. \end{cases} \quad (2.1)$$

Remarks. 1. Notice that Theorem 1 holds not only for d.f. F with a regularly varying tail, but also extends to d.f. with a more general tail behaviour; see [11].

2. It follows from (2.1) that (cf. [16, p. 212])

$$\liminf_{t \rightarrow \infty} \frac{M_t}{L_\alpha\left(\frac{t}{\log \log t}\right)} \left(\frac{\log \log t}{t}\right)^{1/\alpha} = \mu^{-1/\alpha} \quad \text{a.s.} \quad (2.2)$$

We now concentrate on the case $\mu = \infty$.

Theorem 2. *Let*

$$1 - F(x) = \frac{x^{-\alpha}}{L(x)}, \quad 0 \leq \alpha \leq 1$$

and suppose that $\mu = \infty$.

A. *If $0 \leq \alpha < 1$,*

$$\limsup_{t \rightarrow \infty} \frac{M_t}{t} = 1 \quad \text{a.s.}$$

B. (i) *If $\alpha = 1$,*

$$\liminf_{t \rightarrow \infty} \frac{M_t}{L_1^{(1)}\left(\frac{t}{\log \log t}\right)} \frac{\log \log t}{t} = 1 \quad \text{a.s.}$$

with

$$L^{(1)}(x) = \left(L(x) \int_0^x \frac{1}{uL(u)} du \right)^{-1}.$$

(ii) *If $0 < \alpha < 1$,*

$$\liminf_{t \rightarrow \infty} M_t \frac{\log \log t}{t} = \beta(\alpha) \quad \text{a.s.}$$

with $\beta(\alpha)$ the root of the equation

$$\sum_{k=1}^{\infty} \frac{\beta^k}{k!} \frac{\alpha}{k - \alpha} = 1.$$

(iii) *If $\alpha = 0$, $1 - F \in \pi R$,*

$$\frac{1}{x} \int_0^x u dF(u) = O\left(\frac{1 - F(x)}{\log \log \log x}\right),$$

$$x(1 - F(x)) \int_0^x u \left(1 - \log \frac{x}{u}\right) dF(u) = O\left(\left(\int_0^x u dF(u)\right)^2\right) \quad (x \rightarrow \infty), \quad (2.3)$$

then

$$\liminf_{t \rightarrow \infty} \frac{M_t}{a_t} = 1 \quad \text{a.s.}$$

where $a_t^{(1)} \leq a_t \leq a_t^{(2)}$, $(t - a_t^{(1)})Q(a_t^{(1)}) \sim tQ(a_t^{(2)}) \sim \log \log t$,

$$Q(a) = \frac{1}{a} \frac{1 - F(a)}{\frac{1}{a} \int_0^a u \, dF(u)}.$$

Theorem 2.A shows that if $\mu = \infty$ (more precisely if $0 \leq \alpha < 1$), nearly the whole interval $[0, t]$ may be covered by only one renewal period. However we expect this result not to be true in the case $\alpha = 1$. A comparison of (2.2) and Theorem 2.B shows that the lower bound for M_t has a completely different behaviour depending on whether $\mu < \infty$ or not. Indeed, in the first case the parameter dependence can be found in the asymptotic constant while in the second case, the parameter appears in the rate. Also note that in case $0 < \alpha < 1$ the slowly varying function L has no influence.

The case $\alpha = 0$ is very special and we have to concentrate on a special subclass of slowly varying functions. The technical condition in (2.3) is easily checked and is satisfied for a large class of functions $1 - F \in \pi R$. We remark that the bounds of the rate a_t are only given implicitly in (2.3) since they can vary very much depending on the behaviour of $1 - F$, as will be shown by the following examples:

(i) If

$$F(x) = \begin{cases} 0, & x < e, \\ 1 - \exp - \frac{\log x}{\log \log x}, & x \geq e, \end{cases}$$

then

$$\liminf_{t \rightarrow \infty} M_t \left(\frac{\log \log t}{t \log \log \log t} \right) = 1 \quad \text{a.s.}$$

(ii) If

$$F(x) = \begin{cases} 0, & x < 1 \\ 1 - \exp - (\log x)^\gamma, & x \geq 1, \quad 0 < \gamma < 1, \end{cases}$$

then

$$\frac{1 - \gamma}{2 - \gamma} \leq \liminf_{t \rightarrow \infty} t^{-1} M_t \leq \min\{\frac{1}{2}, 1 - \gamma\} \quad \text{a.s.}$$

(iii) If

$$F(x) = \begin{cases} 0, & x < e, \\ 1 - (\log x)^{-\gamma}, & x \geq e, \quad \gamma > 0, \end{cases} \quad (2.4)$$

then

$$\liminf_{t \rightarrow \infty} t^{-1} M_t = \frac{1}{2} \quad \text{a.s.}$$

This last example shows that in case of (2.4), M_t is asymptotically equal to at least half of the whole range t . In order to estimate the remaining part $t - M_t$, we prove the following.

Lemma 3. *Let $M_t^{(1)} \geq M_t^{(2)} \geq \dots$ be the order statistics of the sequence $X_1, X_2, \dots, X_{\tau_t}$, $t - S_{\tau_t}$ and let F be given by (2.4). Then*

$$\lim_{t \rightarrow \infty} \frac{M_t^{(1)} + M_t^{(2)}}{t} = 1 \quad \text{a.s.}$$

Remarks. 1. It is possible to apply Theorems 1 and 2 in order to determine the asymptotic behaviour of

$$\left(\max_{1 \leq i \leq n} X_i \right) S_n^{-1} = M(S_n) \cdot S_n^{-1}.$$

This ratio was investigated among others by Darling [7], Arov and Bobrov [1], and Maller and Resnick [13].

It was shown in [13] that

$$M(S_m) \cdot S_m^{-1} \rightarrow 0 \quad \text{a.s.} \quad \text{if } \mu < \infty \quad (\alpha \geq 1)$$

and

$$M(S_m) \cdot S_m^{-1} \rightarrow 1 \quad \text{a.s.} \quad \text{if } \alpha = 0 \quad \text{and} \quad \int_0^\infty \frac{\int_0^\infty u \, dF(u)}{x(1-F(x))^2} \, dF(x) < \infty.$$

Using Theorem 2.B(iii) we can also deal with the intermediate case $0 < \alpha < 1$: let $S_n \leq t < S_{n+1}$, then

$$t^{-1} M(t) = \begin{cases} t^{-1} M(S_n) & \text{if } S_n \leq t \leq \min\{S_{n+1}, S_n + M(S_n)\}, \\ t^{-1}(t - S_n) & \text{if } \min\{S_{n+1}, S_n + M(S_n)\} \leq t \leq S_{n+1}. \end{cases}$$

Consequently,

$$t^{-1} M(t) \geq (S_n + M(S_n))^{-1} M(S_n) = \frac{S_n^{-1} M(S_n)}{1 + S_n^{-1} M(S_n)} \geq \frac{M(S_n)}{2S_n}. \quad (2.5)$$

Since also

$$\liminf_{t \rightarrow \infty} \frac{M(t)}{t} \leq \liminf_{n \rightarrow \infty} \frac{M(S_n)}{S_n},$$

we have under the conditions of Theorem 2.B(ii) that

$$\liminf_{n \rightarrow \infty} \frac{M(S_n)}{S_n} \log \log S_n = \beta(\alpha) \quad \text{a.s.}$$

2. Several authors (see for example Logan, Mallows, Rice and Shepp [12], McLeish and O'Brien [14] and Csörgő and Horváth [6]) investigated the properties of the sequence

$$S_n^{-1} \left(\sum_{j=1}^n |X_j|^p \right)^{1/p}$$

The case $p = \infty$ corresponds to the case discussed above. It is not clear whether our results can be extended to the case $p < \infty$.

3. The above two remarks suggest that the limit behaviour of $t^{-1}M(t)$ and that of $S_n^{-1}M(S_n)$ agree. However it is not the case. As an example we present the following simple result (without proof): Let F be given by (2.4) then

$$\lim_{n \rightarrow \infty} S_n^{-1}M(S_n) = 1 \quad \text{a.s.}$$

3. Proofs

The proofs of the theorems heavily depend on the following lemmas which are interesting in their own right.

We denote in the sequel $p(t, a)$ as

$$p(t, a) = \begin{cases} P(M_t \leq a) & \text{if } t > a, \\ 1 & \text{if } t \leq a. \end{cases} \quad (3.1)$$

Lemma 4. *Let $p(t, a)$ be defined in (3.1). Then*

$$p(t, a) = \int_0^a p(t-u, a) dF(u).$$

Proof

$$\begin{aligned} P(M_t \leq a) &= E\{P(M_t \leq a | X_1)\} \\ &= E\{P(M_{t-X_1} < a, X_1 < a | X_1)\} \\ &= \int_0^a p(t-u, a) dF(u) \quad \text{if } 0 < a \leq t. \end{aligned}$$

This proves the lemma.

Lemma 5. *Let $a > 0$ and let $y(a)$ be the root of the equation*

$$\int_0^a (\exp uy(a)) dF(u) = 1. \quad (3.2)$$

Then

(i) if $0 < \alpha < 1$,

$$y(a) \sim \frac{\beta(\alpha)}{a} \quad (a \rightarrow \infty)$$

with $\beta(\alpha)$ the root of the equation

$$\sum_{k=1}^{\infty} \frac{\beta^k}{k!} \frac{\alpha}{k-\alpha} = 1;$$

(ii) if $1 \leq \alpha < \infty$,

$$y(a) \sim \frac{1-F(a)}{\int_0^a u \, dF(u)} \quad (a \rightarrow \infty);$$

(iii) if $\alpha = 0$ and the conditions of Theorem 2.B (iii) are satisfied, then

$$y(a) \sim \frac{1}{a} \log \frac{1-F(a)}{\frac{1}{a} \int_0^a u \, dF(u)} \quad (a \rightarrow \infty). \quad (3.3)$$

Proof. (i) Take $a > 0$ such that $y(a) \leq \beta/a$; then from the inequality

$$0 \leq u \left(\frac{\beta}{a} - y(a) \right) \leq \exp \frac{u\beta}{a} - \exp uy(a), \quad 0 \leq u \leq a,$$

we have that

$$\begin{aligned} \left(\frac{\beta}{a} - y(a) \right) \cdot \int_0^a u \, dF(u) &\leq \int_0^a \left(\exp \frac{u\beta}{a} - \exp uy(a) \right) dF(u) = -(1-F(a)) \exp \beta \\ &\quad + \frac{\beta}{a} \int_0^a (1-F(u)) \exp \frac{u\beta}{a} du, \end{aligned} \quad (3.4)$$

where the last equality resulted from an integration by parts and (3.2). It follows from

$$\sum_{k=1}^{\infty} \frac{\beta^k}{k!} \frac{\alpha}{k-\alpha} = 1$$

that β is the unique solution of the integral equation

$$\frac{\exp \beta}{\beta} = \int_0^1 v^{-\alpha} \exp \beta v \, dv.$$

Using this, we get from (3.4) that

$$\left(\frac{\beta}{a} - y(a) \right) \int_0^a u \, dF(u) \leq \beta(1-F(a)) \int_0^1 \left(\frac{1-F(av)}{1-F(a)} - v^{-\alpha} \right) \exp \beta v \, dv.$$

With the regular variation of $1-F$, we have that (see [9])

$$\begin{aligned} 1 - \frac{ay(a)}{\beta} &\leq \frac{a(1-F(a))}{\int_0^a u \, dF(u)} \cdot o(1) \quad (a \rightarrow \infty) \\ &= \frac{1-\alpha}{\alpha} \cdot o(1) \quad (a \rightarrow \infty). \end{aligned}$$

Hence it follows that $\lim ay(a) = \beta$ if the limit is taken over all points a for which $y(a) \leq \beta/a$.

In the same way as above we can show that $\lim ay(a) = \beta$ if the limit is taken over all points a for which $y(a) \geq \beta/a$, and part (i) follows.

(ii) Since

$$e^{uy} - 1 - uy = \int_0^{uy} \int_0^v e^z dz dv \begin{cases} \leq \frac{1}{2} e^{uy} (uy)^2, \\ \geq \frac{1}{2} (uy)^2, \end{cases}$$

it follows from (3.2) that

$$1 = F(a) + y \int_0^a u dF(u) + R \quad (3.5)$$

with

$$\frac{y^2}{2} \int_0^a u^2 dF(u) \leq R \leq \frac{y^2}{2} \int_0^a u^2 e^{uy} dF(u).$$

Moreover, (3.5) implies that

$$y(a) \leq f(a) := \frac{1 - F(a)}{\int_0^a u dF(u)},$$

whence

$$0 \leq 1 - \frac{y(a)}{f(a)} \leq \frac{1}{2(1 - F(a))} f^2(a) e^{af(a)} \int_0^a u^2 dF(u) = V(a).$$

If $\alpha > 1$,

$$\begin{aligned} V(a) &\sim \frac{1}{2\mu^2} (1 - F(a)) \int_0^a u^2 dF(u) \\ &= \frac{1}{2\mu^2} (1 - F(a)) (-a^2(1 - F(a)) + 2 \int_0^a u(1 - F(u)) du) \\ &= \frac{1}{2\mu^2} (-(a(1 - F(a)))^2 + 2a(1 - F(a)) \int_0^a (1 - F(u)) du) \\ &\rightarrow 0 \quad (a \rightarrow \infty). \end{aligned}$$

If $\alpha = 1$,

$$\int_0^a u^2 dF(u) \sim \frac{1}{2} a^2 (1 - F(a)) \quad (a \rightarrow \infty),$$

whence

$$0 \leq 1 - \frac{y(a)}{f(a)} \leq \left(\frac{a(1 - F(a))^2}{\int_0^a u dF(u)} \right).$$

Since

$$\frac{a(1-F(a))}{\int_0^a u dF(u)} = \left\{ \int_0^1 \frac{1-F(av)}{1-F(a)} dv - 1 \right\}^{-1} \rightarrow 0 \quad (a \rightarrow \infty),$$

the proof is completed.

(iii) Define $f(a)$ as the solution of the following equation:

$$\frac{1-F(a)}{1/a \int_0^a u dF(u)} = \sum_{k=1}^{\infty} \frac{(af(a))^k}{k!k}. \quad (3.6)$$

Since $1-F \in \pi V$, the left hand side in (3.6) tends to ∞ , whence also $af(a) \rightarrow \infty$ if $a \rightarrow \infty$. Furthermore,

$$\sum_{k=1}^{\infty} \frac{(af(a))^k}{k!k} = \frac{e^{af(a)}}{af(a)} \left(1 + \frac{1}{af(a)} + O\left(\frac{1}{af(a)}\right)^2 \right) \quad (a \rightarrow \infty), \quad (3.7)$$

so that, with (3.6),

$$\frac{1-F(a)}{1/a \int_0^a u dF(u)} = \frac{e^{af(a)}}{af(a)} \left(1 + \frac{1}{af(a)} + O\left(\frac{1}{af(a)}\right)^2 \right) \quad (a \rightarrow \infty).$$

We now take $z(a)$ such that

$$\frac{1-F(a)}{1/a \int_0^a u dF(u)} = \frac{e^{az(a)}}{az(a)}, \quad a > 0.$$

Then for a sufficiently large we have that $f(a) \leq z(a)$ and by use of the inequality (still for a large enough)

$$\begin{aligned} (az(a) - af(a)) \frac{e^{af(a)}}{af(a)} &\leq \frac{e^{az(a)}}{az(a)} - \frac{e^{af(a)}}{af(a)} \\ &= O\left(\frac{e^{af(a)}}{(af(a))^2}\right) \quad (a \rightarrow \infty), \end{aligned} \quad (3.8)$$

we get that

$$z(a) \sim f(a) \quad (a \rightarrow \infty). \quad (3.9)$$

By taking

$$h(a) = \frac{1}{a} \log \left(\frac{1-F(a)}{1/a \int_0^a u dF(u)} \cdot \log \left(\frac{1-F(a)}{1/a \int_0^a u dF(u)} \right) \right), \quad a > 0,$$

we have that

$$\frac{e^{ah(a)}}{ah(a)} = \frac{1-F(a)}{1/a \int_0^a u dF(u)} \left(1 + \frac{\log \log \frac{1-F(a)}{1/a \int_0^a u dF(u)}}{\log \frac{1-F(a)}{1/a \int_0^a u dF(u)}} \right)^{-1},$$

whence $h(a) \leq z(a)$.

Applying the same inequality as in (3.8) then yields that $z(a) \sim h(a)$, which, combined with (3.9), implies that

$$f(a) \sim \frac{1}{a} \log \frac{1 - F(a)}{1/a \int_0^a u \, dF(u)} \quad (a \rightarrow \infty).$$

We now show in the second part of the proof that

$$y(a) \sim f(a) \quad (a \rightarrow \infty).$$

From integration by parts, we have that

$$\begin{aligned} 1 - \int_0^a (\exp uf(a)) \, dF(u) &= (1 - F(a)) - af(a) \int_0^1 (F(a) - F(av)) \exp vaf(a) \, dv \\ &= (1 - F(a)) - \left(\int_0^a u \, dF(u) \right) f(a) \int_0^1 \left(\log \frac{1}{v} \right) (\exp vaf(a)) \, dv \\ &\quad - af(a) \int_0^1 \left[F(a) - F(av) - \left(\log \frac{1}{v} \right) \left(\frac{1}{a} \int_0^a u \, dF(u) \right) \right] \exp vaf(a) \, dv. \end{aligned}$$

Since

$$af(a) \int_0^1 \left(\log \frac{1}{v} \right) (\exp vaf(a)) \, dv = \sum_{k=1}^{\infty} \frac{(af(a))^k}{k! k}, \quad (3.10)$$

it follows from the choice of $f(a)$ (see (3.6)) and (3.10) that

$$\begin{aligned} 1 - \int_0^a (\exp uf(a)) \, dF(u) &= -af(a) \int_0^1 \left[F(a) - F(av) - \left(\log \frac{1}{v} \right) \left(\frac{1}{a} \int_0^a u \, dF(u) \right) \right] \\ &\quad (\exp vaf(a)) \, dv. \end{aligned} \quad (3.11)$$

Now

$$\begin{aligned} F(a) - F(av) - \left(\log \frac{1}{v} \right) \left(\frac{1}{a} \int_0^a u \, dF(u) \right) &= C(av) - C(a) + \int_1^v (C(au) - C(a)) \frac{du}{u} \end{aligned} \quad (3.12)$$

where $C(a) := 1/a \int_0^a u \, dF(u)$.

Since $1 - F \in \pi R$ we have that $C \in \pi V$ (see (1.9)). It then follows from [2, I, Corollary 3.6] that for every $0 < \varepsilon < 1$ there exists a constant K such that for a large enough,

$$|C(av) - C(a)| \leq K v^{-\varepsilon} \cdot \left| \frac{1}{a} \int_0^a u \, dC(u) \right|$$

uniformly in $v \in (0, 1)$.

Using this in (3.12), we get from (3.11) that for a large enough,

$$\begin{aligned} & \left| 1 - \int_0^a (\exp uf(a)) \, dF(u) \right| \\ & \leq K \left| f(a) \cdot \int_0^a u \, dC(u) \right| \left\{ \int_0^1 v^{-\varepsilon} (\exp vaf(a)) \, dv \right. \\ & \quad \left. + \frac{1}{af(a)} \int_0^1 v^{-1-\varepsilon} (\exp vaf(a) - 1) \, dv \right\} \\ & \leq \frac{2K}{1-\varepsilon} \left| \frac{1}{a} \int_0^a u \, dC(u) \right| \cdot (\exp af(a)). \end{aligned} \quad (3.13)$$

Now take $a > 0$ such that $y(a) \geq f(a)$. Then from the inequality

$$u(y(a) - f(a)) e^{uf(a)} \leq e^{uy(a)} - e^{uf(a)}, \quad 0 \leq u \leq a,$$

we have that

$$\begin{aligned} & \left(\frac{y(a)}{f(a)} - 1 \right) \cdot f(a) \cdot \int_0^a u e^{uf(a)} \, dF(u) \leq \int_0^a (e^{uy(a)} - e^{uf(a)}) \, dF(u) \\ & = 1 - \int_0^a e^{uf(a)} \, dF(u). \end{aligned} \quad (3.14)$$

Using integration by parts, we get that

$$\int_0^a u e^{uf(a)} \, dF(u) = a \int_0^1 (F(a) - F(av)) e^{vaf(a)} (1 + vaf(a)) \, dv$$

and with a similar method as above, we can estimate this integral to find that

$$\int_0^a u e^{uf(a)} \, dF(u) \sim \left(\int_0^a u \, dF(u) \right) \frac{e^{af(a)}}{(af(a))} \quad (a \rightarrow \infty).$$

Combining this estimate with (3.13) and (3.14), it follows from (2.3) that

$$\lim_{a \rightarrow \infty} \frac{y(a)}{f(a)} = 1 \quad (3.15)$$

if the limit is taken over all points a for which $y(a) \geq f(a)$. Take now $a > 0$ such that $y(a) \leq f(a)$. Since

$$a(f(a) - y(a)) \left(\frac{1}{a} \int_0^a u \, dF(u) \right) \leq \int_0^a e^{uf(a)} \, dF(u) - 1,$$

it follows from (2.3), (3.13) and the choice of $f(a)$ that

$$\limsup_{a \rightarrow \infty} a(f(a) - y(a)) < \infty. \quad (3.16)$$

We now apply the same inequality as in (3.14) with $f(a)$ and $y(a)$ interchanged, then by (3.13) and (3.16),

$$\lim_{a \rightarrow \infty} \frac{y(a)}{f(a)} = 1$$

if the limit is taken over all points a for which $y(a) \leq f(a)$. Combining this with (3.15) proves the lemma.

Lemma 6. *Let $p(t, a)$ and $y(a)$ be defined as in (3.1) and (3.2). Then for every $\alpha \geq 0$, there exist absolute constants $C_1(\alpha)$, $C_2(\alpha) > 0$ such that*

$$\begin{aligned} p(t, a) &= C_\alpha(t, \alpha) \exp(-ty(a)) && \text{if } \alpha > 0, \\ C_1(0) \exp(-ty(a)) &\leq p(t, a) \leq C_2(0) \exp(-(t-a)y(a)) && \text{if } \alpha = 0, \end{aligned} \quad (3.17)$$

and

$$C_1(\alpha) \leq C_\alpha(t, a) \leq C_2(\alpha).$$

Proof. We first consider the case $\alpha > 0$. In this case Lemma 5 implies the existence of two constants $L_1(\alpha) \geq 0$ and $L_2(\alpha) > 0$ such that

$$L_1(\alpha) \leq ay(a) \leq L_2(\alpha).$$

Since, for $t \leq a$,

$$\begin{aligned} 1 &= p(t, a) \\ &= C_\alpha(t, a) \exp(-ty(a)) \begin{cases} \geq C_\alpha(t, a) \exp(-L_2(\alpha)), \\ \leq C_\alpha(t, a), \end{cases} \end{aligned}$$

it follows that

$$1 \leq C(t, a) \leq \exp L_2(\alpha). \quad (3.18)$$

We now show that (3.18) also holds for $t > a$. Suppose that this is not the case, then there exists a smallest $t = t_0 > a$ for which $C(t_0, a) = e^{L_2(\alpha)}$ (or $C(t_0, a) = 1$). By

Lemma 4,

$$\begin{aligned}
 p(t_0, a) &= C(t_0, a) \exp(-t_0 y(a)) \\
 &= \int_0^a C(t_0 - u, a) \cdot (\exp(-(t_0 - u)y(a))) dF(u) \\
 &= \exp(-t_0 y(a)) \int_0^a C(t_0 - u, a) (\exp uy(a)) dF(u) \\
 &< e^{L_2(\alpha)} \exp(-t_0 y(a)) \int_0^a (\exp uy(a)) dF(u) \\
 &= e^{L_2(\alpha)} \exp(-t_0 y(a)),
 \end{aligned}$$

whence

$$C(t_0, a) < e^{L_2(\alpha)},$$

giving a contradiction. Hence (3.18) holds for all t and a .

The proof of the case $\alpha = 0$ is completely similar. The details are left to the reader.

Proof of Theorem 2.A. Let

$$n_i(i: 1, 2, \dots) = \begin{cases} (2^i)^{1/\alpha} L_\alpha(2^i) & \text{if } 0 < \alpha < 1, \\ L^{\text{inv}}(2^i) & \text{if } \alpha = 0, \end{cases}$$

where L^{inv} is the inverse of L . Notice that L^{inv} is well defined, since if $\alpha = 0$, L is monotone.

Further, let

$$n_i^* := \frac{1}{1 - F(n_i)}, \quad i = 1, 2, \dots$$

Define the events ($i = 1, 2, \dots$)

$$\begin{aligned}
 A_i &= A_i(\varepsilon) = \left\{ \max_{1 \leq j \leq n_i^*} X_j \geq n_i, \sum_{j=1}^{n_i^*} X_j - \max_{1 \leq j \leq n_i^*} X_i \leq \varepsilon n_i \right\}, \\
 B_i &= \left\{ \max_{n_i^* \leq j < n_{i+1}^*} X_j \geq n_{i+1}, \sum_{j=n_i^*}^{n_{i+1}^*-1} X_j - \max_{n_i^* \leq j < n_{i+1}^*} X_j \leq \varepsilon n_{i+1} \right\}
 \end{aligned}$$

and

$$B_i^{(k)} = \left\{ X_k \geq n_{i+1}, \sum_{\substack{j=n_i^* \\ j \neq k}}^{n_{i+1}^*-1} X_j \leq \varepsilon n_{i+1} \right\}, \quad n_i^* \leq k < n_{i+1}^*.$$

If $0 < \alpha < 1$, it follows from Feller [10] that

$$P(B_i^{(k)}) \sim \frac{\Lambda_\alpha(\varepsilon)}{n_{i+1}^*} \quad (i \rightarrow \infty) \quad (3.19)$$

where Λ_α is a stable law of index α .

If $\alpha = 0$, we have from Darling [7] that

$$P(B_i^{(k)}) \sim \frac{e^{-1/2}}{n_{i+1}^*} \quad (i \rightarrow \infty). \quad (3.20)$$

Since the events $B_i^{(k)}$ ($k = n_i^*, \dots, n_{i+1}^* - 1$) are disjoint (provided that $0 < \varepsilon < 1$), and since

$$\sum_{k=n_i^*}^{n_{i+1}^*-1} B_i^{(k)} = B_i,$$

it follows from (3.19) and (3.20) that

$$P(B_i) \rightarrow C > 0 \quad \text{if } 0 \leq \alpha < 1.$$

Since furthermore the events $(B_i)_{i=1}^\infty$ are mutually independent, the Borel-Cantelli Lemma implies that the events B_i occur infinitely often (for any $0 < \varepsilon < 1$), which in turn implies that the events A_i occur infinitely often.

Let $N_1 = N_1(\omega) < N_2 = N_2(\omega) < \dots$ be a random sequence of integers such that A_{N_i} occurs for $i = 1, 2, \dots$. Define $t_i = S_{N_i^*}$ where $N_i^* = (1 - F(N_i))^{-1}$.

Then

$$t_i \leq \varepsilon N_i + \max_{1 \leq j \leq N_i^*} X_j$$

and

$$M_{t_i} = \max_{1 \leq j \leq N_i^*} X_j.$$

Consequently $t_i^{-1} M_{t_i} \geq (1 + \varepsilon)^{-1}$ for any $1 > \varepsilon > 0$. Since $t^{-1} M_t \leq 1$ ($t > 0$), the proof is finished.

Proof of Theorem 2.B. Take $\varepsilon > 0$. Choose $a_\alpha(t)$ ($b_\alpha(t)$) as the solution of

$$y(a_\alpha(t)) = (1 + \varepsilon) \frac{\log \log t}{t} \quad \left(y(b_\alpha(t)) = (1 - \varepsilon) \frac{\log \log t}{t} \right)$$

if $\alpha > 0$ and let $a_0(t)$ ($b_0(t)$) be defined by

$$\begin{aligned} (t - a_0(t)) y(a_0(t)) &= (1 + \varepsilon) \log \log t, \\ t y(b_0(t)) &= (1 - \varepsilon) \log \log t. \end{aligned}$$

Now let f be defined as

$$f(a) = \begin{cases} \frac{1}{a} \log \frac{1 - F(a)}{\frac{1}{a} \int_0^a u \, dF(u)} & \text{if } \alpha = 0, \\ \frac{\beta(\alpha)}{a} & \text{if } 0 < \alpha < 1, \\ \frac{1 - F(a)}{\int_0^a u \, dF(u)} & \text{if } \alpha = 1, \end{cases} \quad (3.21)$$

$$f(a) = \begin{cases} \frac{\beta(\alpha)}{a} & \text{if } 0 < \alpha < 1, \end{cases} \quad (3.22)$$

$$f(a) = \begin{cases} \frac{1 - F(a)}{\int_0^a u \, dF(u)} & \text{if } \alpha = 1, \end{cases} \quad (3.23)$$

and take $c_\alpha(t)$ such that

$$(1 + \varepsilon)^{-1} y(a_\alpha(t)) = f(c_\alpha(t)) = (1 - \varepsilon)^{-1} y(b_\alpha(t)) \quad \text{for } 0 \leq \alpha \leq 1.$$

It then follows from Lemma 5 that

$$\frac{f(a_\alpha(t))}{f(c_\alpha(t))} = \frac{f(a_\alpha(t))}{y(a_\alpha(t))} (1 + \varepsilon) \rightarrow (1 + \varepsilon) \quad (t \rightarrow \infty).$$

Since f in (3.21)–(3.23) is regularly varying with a non-zero index, it follows from [9, Corollary 1.2.1] that

$$\lim_{t \rightarrow \infty} \frac{a_\alpha(t)}{c_\alpha(t)} = \begin{cases} (1 + \varepsilon)^{-1} & \text{if } 0 \leq \alpha < 1, \\ (1 + \varepsilon)^{-\alpha} & \text{if } \alpha = 1. \end{cases} \quad (3.24)$$

In the same way we can show that

$$\lim_{t \rightarrow \infty} \frac{b_\alpha(t)}{c_\alpha(t)} = \begin{cases} (1 - \varepsilon)^{-1} & \text{if } 0 \leq \alpha < 1 \\ (1 - \varepsilon)^{-\alpha} & \text{if } \alpha = 1. \end{cases} \quad (3.25)$$

By Lemma 6 and the choice of $a_\alpha(t)$, we have that, for $0 \leq \alpha \leq 1$,

$$p(t, a_\alpha(t)) \leq C_2(\alpha) (\log t)^{-(1+\varepsilon)}.$$

Now take

$$t_k = \theta^k, \quad \theta > 1, \quad k = 1, 2, \dots;$$

then

$$P(M_{t_k} < a_\alpha(t_k)) \leq C_2(\alpha) (k \log \theta)^{-(1+\varepsilon)}$$

whence by the Borel-Cantelli Lemma,

$$M_{t_k} \geq a_\alpha(t_k) \quad \text{a.s.}$$

for all but finitely many k .

Since M_t is a decreasing function of t , for any $t_k \leq t < t_{k+1}$, we obtain that $M_t \geq a_\alpha(t_k)$. Choosing θ sufficiently close to 1, we get that

$$M_t \geq (1 - \varepsilon) a_\alpha(t) \quad \text{a.s.}$$

if t is big enough.

Now using (3.24) and letting $\varepsilon \rightarrow 0$ gives

$$\liminf_{t \rightarrow \infty} \frac{M_t}{c_\alpha(t)} \geq 1 \quad \text{a.s.} \quad (3.26)$$

We now turn to the proof of the converse in (3.26) and we show that

$$\liminf_{t \rightarrow \infty} \frac{M_t}{c_\alpha(t)} \leq 1 \quad \text{a.s.} \quad (3.27)$$

Let $\alpha > 0$, $u_k = \exp(k \log k)$ and let m_k be the smallest integer for which

$$S_{m_k} \geq u_k.$$

Observe that

$$u_k = o(b_\alpha(u_{k+1})) \quad (k \rightarrow \infty)$$

and

$$\begin{aligned} P(S_{m_k} \geq \varepsilon b_\alpha(u_{k+1})) &\leq P(X_{m_k} \geq \varepsilon b_\alpha(u_{k+1}) - u_k) \\ &\leq P(X_{m_k} \geq \frac{\varepsilon}{2} b_\alpha(u_{k+1})) \\ &\leq P\left(X_1 \geq \frac{\varepsilon}{2} b_\alpha(u_{k+1}) \mid X_1 \geq u_k\right) \\ &\leq P(X_1 \geq \frac{\varepsilon}{2} \frac{b_\alpha(u_{k+1})}{P(X_1 \geq u_k)}). \end{aligned}$$

Using (3.25) and the definition of $c_\alpha(t)$, it follows with an application of the Borel-Cantelli Lemma that for every $\varepsilon > 0$,

$$S_{m_k} \leq \varepsilon \cdot b_\alpha(u_{k+1}) \quad \text{a.s.} \quad (3.28)$$

for all but finitely many k .

Let

$$M(S_{m_k}, u_{k+1}) = \max\{X_{m_{k+1}}, X_{m_{k+2}}, \dots, X_{\tau_{u_{k+1}}}, u_{k+1} - S_{\tau_{u_{k+1}}}\},$$

then by Lemma 6 and the choice of $b_\alpha(t)$,

$$P(M(S_{m_k}, u_{k+1}) < b_\alpha(u_{k+1})) \geq c(\alpha)(\log u_{k+1})^{-(1-\varepsilon)}$$

where $c(\alpha)$ is some absolute constant.

Hence by the Borel-Cantelli Lemma, we have that

$$M(S_{m_k}, u_{k+1}) \leq b_\alpha(u_{k+1}) \quad \text{i.o. a.s.}$$

which together with (3.28) and (3.25) implies (3.27). In case $\alpha = 0$, let $u_k = L^{\text{inv}}(\exp(k \log k))$ and we get the result similarly.

Proof of Lemma 3. Let

$$T = T(t) = [2(\log t)^\gamma \log \log t];$$

then

$$P(\max(X_1, \dots, X_T) < 2t) = (F(2t))^T = \left(1 - \frac{1}{(\log 2t)^\gamma}\right)^T \sim \frac{1}{(\log t)^2} (t \rightarrow \infty).$$

Now take $t_k = 2^k$; then

$$\max(X_1, \dots, X_{T(t_k)}) \geq 2t_k \quad \text{a.s.}$$

if k is big enough. Further let $t_k \leq t < t_{k+1}$; then

$$\max(X_1, \dots, X_{T(t)}) \geq \max(X_1, \dots, W_{T(t_k)}) \geq 2t_k \geq t.$$

Hence

$$\max(X_1, \dots, X_{T(t)}) \geq t \quad \text{a.s.}$$

if t is big enough. This in its turn implies that

$$\tau_t \leq T. \tag{3.29}$$

We now show that

$$\xi_t := \#\left\{i: i \leq T(t), \alpha(t) = \frac{t}{(\log t)^{2\gamma}} \leq X_i \leq t = \beta(t)\right\} \leq 1 \quad \text{a.s.}$$

if t is big enough. Let

$$\xi_t^* := \#\{i: i \leq T(2t), \alpha(t) \leq X_i \leq \beta(2t)\}.$$

Since

$$\begin{aligned} P_t &:= P(\alpha(t) \leq X_i \leq \beta(2t)) = \left(\log\left(\frac{t}{(\log t)^{2\gamma}}\right)\right)^{-\gamma} - (\log 2t)^{-\gamma} \\ &\sim 2\gamma^2 \frac{\log \log t}{(\log t)^{\gamma+1}} \quad (t \rightarrow \infty), \end{aligned}$$

we have that

$$\begin{aligned} P(\xi_t^* > 1) &= 1 - (1 - P_t)^{T(2t)} - T(2t)(1 - P_t)^{T(2t)-1} \cdot P_t \\ &\sim \frac{(T(2t)P_t)^2}{2} \\ &\sim 8\gamma^4 \frac{(\log \log t)^4}{(\log t)^2} \quad (t \rightarrow \infty). \end{aligned}$$

Let $t_k = 2^k$. Then $\xi_{t_k}^* \leq 1$ a.s. if k is big enough. Now take

$$t_k \leq t \leq t_{k+1}.$$

Since

$$T(t) \leq T(2t_k)$$

and

$$\alpha(t_k) \leq \alpha(t) < \beta(t) \leq \beta(2t_k),$$

we obtain that

$$\xi_t \leq \xi_{t_k}^* \leq 1.$$

Since $\tau_t \leq T(t)$ (see (3.29)) we have that $S_{T(t)} \geq t$. Clearly $M_t^{(2)} < t$ and the sum of those elements of the sequence $X_1, X_2, \dots, X_{\tau_t}$, $t - S_{\tau_t}$ which are smaller than $\alpha(t)$ is $o(t)$. Hence the fact that $\xi_t \leq 1$ a.s. implies the lemma.

Acknowledgement

Research done while the first named author visited Catholic University of Leuven, Department of Mathematics (Aug 1, 1986–Aug 31, 1986). He wishes to express his best thanks to the Department and especially to Prof. J. Teugels for their kind hospitality. The authors are indebted to Prof. G. O'Brien who suggested Remarks 1 and 2 following Lemma 3, and to the referee for many comments which lead to an improvement in presentation.

References

- [1] D.Z. Arov and A.A. Boborov, The extreme terms of a sample and their role in the sum of independent variables, *Theory Prob. Appl.* 5 (1960) 377–395.
- [2] N.J. Bingham and C.M. Goldie, Extensions of regular variation I and II, *Proc. Lond. Math. Soc.* (e) 44 (1983) 473–534.
- [3] N.H. Bingham and J.L. Teugels, Duality for regularly varying functions, *Quarterly J. Math.* 26 (1475) 333–353.
- [4] K.L. Chung and P. Erdős, On the application of the Borel–Cantelli lemma, *Trans. Am. Math. Soc.* (1952) 179–186.
- [5] E. Csáki, P. Erdős and P. Révész, On the length of the longest excursion, *Z. Wehrsch. Verw. Geb.* 68 (1985) 365–382.
- [6] M. Csörgő and L. Horváth, Asymptotic representations of self-normalized sums, *Prob. Math. Stat.* (Wroclaw), in press.
- [7] D.A. Darling, The influence of the maximum term in addition of independent random variables, *Trans. Amer. Math. Soc.* 73 (1952) 95–107.
- [8] N.G. de Bruyn, Pairs of slowly oscillating functions occurring in asymptotic problems concerning the Laplace-transformation, *Nieuw Archief Wiskunde* 7 (1959) 20–26.

- [9] L. de Haan, On regular variation and its applications to the weak convergence of sample extremes, Math. Center Tracts (Amsterdam, 1975).
- [10] W. Feller, An introduction to probability theory and its applications, vol. II (Wiley and Sons, New York, 1971).
- [11] J. Galambos, The asymptotic theory of extreme order statistics (Wiley and Sons, New York, 1978).
- [12] B.F. Logan, C.L. Mallows, S.O. Rice and L.A. Shepp, Limit distributions of self normalized sums, Ann. Probability 1 (1973) 788–809.
- [13] R.A. Maller and S.I. Resnick, Limiting behaviour of sums and the term of maximum modulus, Proc. Lond. Math. Soc. (3) 49 (1984) 385–422.
- [14] D.L. McLeish and G.L. O'Brien, The expected ratio of the sum of squares to the square of the sum, Ann. Probability 10 (1982) 1019–1028.
- [15] E. Omey and E. Willekens, π -variation with remainder, to appear in J. Lond. Math. Soc. (1987).
- [16] M.J. Wichura, On the functional form of the law of the iterated logarithm for the partial maxima of independent identically distributed random variables, Ann. Prob. 2 (1974) 202–230.